

Nutritional Benefits of Black Soldier Fly Larvae Oil in Animal Feed

Rudy Agung Nugroho*, Retno Aryani, Esti Handayani Hardi, Hetty Manurung, Rudianto Rudianto

Faculty of Fisheries and Marine Science, Mulawarman Universitas. Jl. Barong Tongkok No 4 Gn Kelua, Samarinda 75123, Kalimantan Timur, Indonesia

Corresponding e-mail : rudyagung.nugroho@fmipa.unmul.ac.id

Received: 12 January 2024, Revised: 19 March 2024, Available Online: 26 May 2024

Abstract – Black Soldier Fly Larvae (BSFL) oil is currently an insect oil that potentially replaces fish oils in animal feed. However, there are limited information on its benefits and nutritional comparison of BSFL oil to fish oil. Present study was to evaluate the benefits of BSFL oil in animal feed and compare the BSFL to Fish oil. Collected data of BSFL oil in the animal feed and its physiological effects was obtained from several articles of reputable journal and compiled into a table. Meanwhile, BSFL was reared and extract their oil to determine the fatty acid composition and compare to fish oil. The current study indicated that BSFL oil can be used in various animal feeds, such as aquafeed, broiler feed, and pets. The BSFL oil which is included in the feed showed an improvement in growth and other physiological responses of animals. Meanwhile, in comparison to fish oil, BSFL oil has a similar composition of fatty acids such as Omega 3 and Omega 6. Eicosapentaenoic acid (EPA; 20:5n3) is detected in the BSFL oil, though the quantity is much lower than in fish oil. Further, the BSFL oil also contains lauric acid. In conclusion, BSFL oil can be an alternative feed ingredient to substitute fish oil and has beneficial to the growth and physiological responses of the animal

Keywords: BSFL, Feed, Feish, Oil

INTRODUCTION

Insects have been utilized both as a substitute for traditional oil sources, such as fish oil (FO) and soybean oil (SBO) as a source of protein (Devi et al., 2022; Kipkoech et al., 2023; Nugroho & Nur, 2018), but they are also becoming value-added feed resources for the animal feed diets (Heuel et al., 2021; Kierończyk et al., 2022; Y. B. Kim et al., 2020). During their development, black soldier fly larvae, also known as BSFL (*Hermetia illucens* L.), are able to transform the huge amounts of organic substrates found in food or animal wastes into edible protein and fats/oil (Gligorescu et al., 2020; Lim et al., 2019; Liu et al., 2018). This makes them a popular bio recycling organism. According to Patterson et al. (2021); Wang and Shelomi (2017) the BSFL has more than forty percent protein that is abundant in essential amino acids and more than twenty-eight percent lipids. Meanwhile, the BSFL oil has a high concentration of medium-chain fatty acids, such as palmitic acid (C16:0), which is analogous to coconut oil (Abduh et al., 2017;

Ushakova et al., 2016). The coconut oil is the only plant-origin oil where about 50% of the fatty acid composition is lauric acid (C12:0) (Nasir et al., 2018) which is also found in the BSFL oil (Lumanlan et al., 2022).

Due to their preference for energy consumption, Wang et al. (2015) found that medium-chain fatty acids may be advantageous in reducing abdominal fat. This is because medium-chain fatty acids consume less energy than long-chain saturated or unsaturated fatty acids. In addition, studies have demonstrated that lauric acid and other medium-chain fatty acids, like it, have antibacterial effects on the bacteria that live in the gut (Schiavone et al., 2017; Zeitz et al., 2015). Therefore, insect oils that are rich in lauric acid may have an influence on the growth performance and intestinal health of broiler chickens, who are known for their rapid rate of development (A Schiavone et al., 2018).

According to Khatun et al. (2018); Taulescu et al. (2010), the amount of fatty acids found in grill feed has an effect on the levels of those

acids found in meats and internal organs. This has direct implications for fat accumulation and metabolism. Meanwhile, Schiavone et al. (2017) discovered that substituting BSFL oil for SBO affected the fatty acid profile of broiler chicks. It was suggested by Kierończyk et al. (2018) that insect oil, such as that extracted from *Tenebrio molitor* and *Zophobas morio*, might be utilized as a replacement for soybean oil in the feed of broiler chickens without affecting either the growth performance or the nutritional digestibility of the chickens. In addition, Li et al. (2017) discovered that consumption of BSFL oil increased the amount of omega-3 fatty acid deposition in the muscles of juvenile carp while simultaneously reducing the amount of intraperitoneal fat buildup.

On the other hand, FO is also a pivotal part of the animal diet. Fish oil is a rich source of omega-3 fatty acids, which are essential nutrients that cannot be produced by animals themselves and must be obtained from their diet. Omega-3 fatty acids play a vital role in many physiological processes in animals, including brain function, immune system regulation, and cardiovascular health. In addition, these fatty acids are important for the development and maintenance of healthy skin and coat in animals. Therefore, including FO in the diet of animals can provide important growth and health benefits.

Though some researchs have been performed in the effects of the BSFL oil and FO in animal diet, However, recent data about the BSFL oil as a partial or total FO/SBO replacement and its physiological effects is still limited. Current research is to compile the physiological effects of the BSFL oil in the diet of fish and poultry animals. The comparison between the BSFL oil and FO are also evaluated in order to understand the potential use in the future feed preparation.

MATERIALS AND METHODS

1. Data comparison for BSFL oil in animal feed

All data regarding the use of BSFL oil in animal feed and their physiological effects was collected from the recent published articles (2018-2023), from various publisher. The information of the BSFL oil in the animal feed and its physiological effects was compiled into a table.

2. BSF rearing.

To compare the fatty acid content of BSFL oil versus FO, several steps were prepared, from BSFL rearing to BSFL oil analysis. In BSFL rearing, a number of BSFL (Average initial weight of 0.05 ± 0.004 g; Initial length of 12.43 ± 0.25 mm) was purchased from a BSF farmer, Samarinda City, East Kalimantan, Indonesia. The BSFL were reared from third-generation flies fed with fermented Palm Kernel Meal (PKM). The BSFL were reared in a plastic box and maintained at temperature of 28.5°C , for 17 days.

3. BSFL oil preparation

At the day 17th, the BSF was in the prepupal stage. All prepupal BSF were collected and dried (60°C) for 48 h, and crushed, to make a meal. The BSFL meal and its oil were separated using an oil extractor (RG-307, Jiangsu, China). The BSFL oil was analyzed with GCMS for fatty acid composition and compared to fish oil.

RESULTS AND DISCUSSION

1. BSFL oil in animal feed

The black soldier fly larvae, often known as BSFL, are a highly sustainable insect candidate species because of their potential for use as a component in animal feed and their capacity to generate revenue. The introduction of insect meals into animal diets

has been shown in a number of studies to have the potential to improve animal survival rates, antioxidant capacities, and immunological responses. Present study

found that various inclusion of BSFL oil in the diet of several animals and its physiological effects (Table 1).

Table 1. Partial and total inclusion/substitution of black soldier fly larvae (BSFL) oil in the diet of several animals and its physiological effects

Species	Suggested Inclusion/substitution	Physiological effects	References
Rainbow trout (<i>Oncorhynchus mykiss</i>)	Potentially be superior to 10%	Posterior intestine did not affect Blood glucose and BSFL oil addition concentration has a negative correlation Several antihyperglycemic effects in fasted rainbow trout. The digestibility of hydroxyproline was higher in feed with BSFL oil compare to reference feed	Dumas et al. (2018)
Rainbow trout (<i>Oncorhynchus mykiss</i>)	16% in the diet	Some immunological benefits Hepar or gut histology did not affect by BSFL oil. Upregulated of kidney interleukin-8, 41 tumor necrosis factor and IRF-1 Significantly higher serum peroxidase activity A substantial increase in lysozyme activity	Kumar et al. (2021)
Siberian sturgeon	30% inclusion	Positive effects on growth performance	Rawski et al. (2020)
Broilers	50% and 100% (Part and full replacement of Soybean oil)	No negative effect on the cecal microbiota of broilers. No minor affected on the Growth statues	B. Kim et al. (2020) B. Kim et al. (2020)
Broiler	25; 50; 75; 100% of BSFL fat (Part and full replacement of Soybean oil)	Several growth performances, length and weight of organ, nutritional absorption coefficients in the ileum did not influence by BSFL oil inclusion	Kierończyk et al. (2020)
Hens	20 g/kg of diet	The quality of the eggs was not affected significantly in terms of shell strength, egg composition, albumen percentage, or any of the major characteristics of the yolk.	Heuel et al. (2021)

Species	Suggested Inclusion/substitution	Physiological effects	References
Hens	Up to 4.5% of soybean oil substitution	Body weight, production of egg, Neither the feed conversion ratio nor the egg weight were affected by the BSFL oil. Eggs produced by hens given BSFL oil had a deeper yellow yolk colour than eggs laid by hens used as controls.	Patterson et al. (2021)
Turkey	50% and 100% (Part and full replacement of Soybean oil)	Growth performance was not influence, nutrient digestibility, gastro-intestinal tractus morphology, or the quality of the muscles of the chest and thighs. Reduced trypsin activity (Total replacement), but this had no effect on digestibility. full change of soybean oil reduced proliferation of potentially pathogenic bacteria (<i>Enterobacteriaceae</i> spp). Decreased levels of IL-6 (Total replacement), while part change reduced the TNF- α concentration.	Sypniewski et al. (2020)
Turkey	50% and 100% (Partial and total replacement of Soybean oil)	Lipase and amylase activity was improved. Liver health status was enhanced by decreased cholesterol and triglyceride concentration. Positively inhibited Bacteroides-Prevotella growth in the cecal digesta. Modulates the immune system by decreasing the level of IL6	Kierończyk et al. (2022)

As can be seen that the BSFL oil has been included and used as inclusion/substitution in various animal diet and has several physiological effects. In fish diet, the BSFL oil inclusion/substitution affected on the growth and immune parameters of the fish. In contrast, Li et al. (2017) observed that no change in growth parameter of the fish when SBO was substituted with BSFL oil from 0 to 100% in *Cyprinus carpio* diets, but did discover a rise in intraperitoneal fat, possibly owing to a considerable elevation of peroxisome proliferator-activated receptor alpha

expression. Meanwhile, Dumas et al. (2018) found no deleterious effects of dietary BSFL oil incorporation levels of 2.5, 5.0, or 10.0% on rainbow trout growth, feeding efficiency, or gut histomorphology, and felt that greater inclusion levels were achievable. Furthermore, the BSFL oil had no impact on the expression of genes involved for inflammation in the gut or any signs of increased fat accumulation in the liver of fish (Kumar et al., 2021). In addition, the BSFL oil may totally replace fish oil or soybean oil in rainbow trout diets while also offering some health advantages, such as enhance the

oxidative state and diminish the expression of pro-inflammatory genes in fish.

Meanwhile, the BSFL oil has a valuable physiological effect on some poultry animals such as hens and turkeys. It is revealed that replacement either partially or totally had positive effects on the health statues of those animals, but no growth effects. The BSFL oil has a genuine position as nutritional supplements for commercial poultry animals. According to Patterson et al. (2021), the oil extracted from BSFL has enough energy (7840 kcal/kg) and linoleic acid (15.29%) to easily replace soybean oil in poultry animal diet. With higher levels of dietary inclusion, for example in hens, both hen performance and egg quality were maintained, and melanin-lauric acid complexes in the amber hue of the BSFL oil are likely transferred to boost egg yolk color.

Furthermore, it is generally understood that the liver's quality, as the organ involved in lipid metabolism, is critical in determining the feasibility of fat/oil source adoption in animal diets. The BSFL oil may be used safely for partial or entire replacement of soybean oil with no changes in biochemical blood characteristics such as AST and ALT levels (Kierończyk et al., 2022). Previous findings of Schiavone et al. (2017); A. Schiavone et al. (2018), in which partial (50%) or entire inclusion of BSFL oil also mentioned that BSFL oil had no effect on AST and ALT blood concentrations in broilers compared to the soybean oil group (SO). Nonetheless, the partial substitution of BSFL oil, as opposed to the total replacement, had the lowest AST score, indicating that partial replacement is more helpful than whole replacement in terms of liver function.

It is hypothesized that changing the kind of oil used in turkey meals from soybean

oil to BSFL oil would have a major impact on the selected microbiota populations in the ceca, which would have led to a reduction in the growth of the *Bacteroides-Prevotella* cluster in the digesta. It is also widely recognized that dietary fats that include high amounts of SFAs from both plant and animal sources, such as palm kernel fatty acid distillers (PKFD) and bovine tallow, may enhance the abundance of potentially harmful bacteria, such as *Bacteroides* and *Clostridium perfringens*. This is because SFAs are a kind of saturated fatty acid that may come from either plants or animals. (Józefiak et al., 2016; Knarreborg et al., 2002). On the other hand, oils that are high in fatty acids (FAs), such as oleic (C18:1), linoleic (C18:2), and linolenic (C18:3) acids, have antibacterial properties (Yoon et al., 2018). The relatively large amount of lauric acid in BSFL oil may have an effect on the microbiota composition, despite the fact that SFAs are the primary components of the FA profile in this oil. The antibacterial, antiviral, and antiparasitic effects that lauric acid has are well known and generally recognized (Dayrit, 2015; Ushakova et al., 2016; Zeitz et al., 2015).

2. BSFL oil versus Fish oil (FO)

Black soldier fly larvae oil and FO are both sources of omega-3 fatty acids, which are essential for good health for animal and abundance with some important fatty acid ingredients (Table 2). The BSFL oil is a relatively new and emerging source of oil that is gaining interest for its potential health benefits.

Table 2. Fatty acid comparison between black soldier fly larvae oil and fish oil

Fatty Acid	BSFL oil	Fish oil
Lauric acid (C12:0)	42.42	nd
Myristoleic Acid (C14:1)	0.23	1.97
<i>cis</i> -10-Pentadecenoic Acid (C15:1)	3.92	0.28
Palmitoleic Acid (C16:1)	0.14	2.45
Heptadecanoic Acid (C17:0)	0.03	6.46
Linoneic Acid (C18:2n6c)	7.15	1.61
Linoleladic Acid (C18:2n6t)	0.77	0.84
γ -Linolenic Acid (C18:3n6)	0.09	0.80
α -Linolenic Acid (C18:3n3)	0.07	2.64
Arachidic Acid (C20:0)	0.96	43.70
<i>cis</i> -11-Eicosanoic Acid (C20:1n9)	0.01	3.84
<i>cis</i> -5,8,11,14,17-Eicosapentaeonic Acid (C20:5n3)	0.04	1.34
Heneicosanoic Acid (C21:0)	0.00	2.52
Behenic Acid (C22:0)	0.01	1.66
Eruchic Acid (C22:1n9)	0.01	22.75
Tricosanoic Acid (C23:0)	0.23	1.97

Noted: W/W%= grams per 100 grams of sample. Results are expressed on a dry weight basis. nd = not detected.

There are some characteristics of the BSFL oil. The BSFL oil is a good source of omega-3 fatty acids, particularly alpha-linolenic acid (ALA). The ALA from BSFL oil is a precursor to other forms of omega-3s that are important for brain and heart health, such as EPA and DHA. Meanwhile, the BSFL oil is also high in lauric acid, which is a medium-chain fatty acid that has been linked to numerous health benefits, including improved cholesterol levels. The BSFL oil contains antioxidants such as tocopherols, which may help protect against oxidative stress and inflammation in the body. The high amount of lauric acid (C12:0) in the BSFL oil is one of its distinguishing characteristics (Oonincx et al., 2015). When animals eat lauric acid, it is promptly oxidized rather than retained in the liver, resulting in decreased feed intake (Belghit et al., 2019).

However, there are some differences between the BSFL oil and FO. Fish oil typically contains more omega-3 fatty acids

than black soldier fly larvae oil (St-Hilaire et al., 2007). This is because fish accumulate omega-3s by consuming algae, which is rich in these fatty acids. The BSFL oil, on the other hand, contains a different type of omega-3 called alpha-linolenic acid (ALA), which is converted by the body into other forms of omega-3s such as EPA and DHA (Lozica et al., 2022). The EPA and DHA are more easily absorbed by the body than ALA, which means that fish oil may be more effective at increasing levels of these important omega-3s in the body. Fish oil contains other important nutrients such as vitamin D and astaxanthin, which have additional health benefits. The BSFL oil contains some antioxidants and vitamins (Lu et al., 2022), but not at the same levels as FO. Further, the BSFL oil is considered more sustainable than FO, as it can be produced using food waste and does not require wild fish populations to be harvested (Hossain et al., 2023). The BSFL oil is also considered to

be a sustainable source of oil because it can be produced using food waste and does not require the harvesting of wild fish populations. In addition, the BSFL oil has been shown to be stable at room temperature for up to six months, which is longer than some other vegetable oils.

CONCLUSION

There are various physiological effects of animal, such as in fish and poultry when the BSFL oil is included either partially or totally replacement in animal diet. Both FO and BSFL oil can be good sources of omega-3 fatty acids. The BSFL oil is a promising new source of oil that is rich in omega-3s, lauric acid, and antioxidants.

Further research is needed to fully understand its health benefits and potential uses of BSFL oil in various animal feed and its gene regulation.

ACKNOWLEDGEMENTS

All Authors grateful thank to Ministry of Education and Culture, Research and Technology for supporting with funding “Penelitian Dasar Unggulan Perguruan Tinggi”, contract Number: 031/E5/PG.02.00.PL/2023 and 300/UN17.L1/HK/2023 Financial Year 2023. Thank also extended to Department of Biology Faculty of Mathematics and Natural Sciences, Mulawarman University, Samarinda, East Kalimantan, Indonesia.

REFERENCES

- Abduh, M. Y., Jamilah, M., Istiandari, P., Manurung, S., & Manurung, R. (2017). Bioconversion of rubber seeds to produce protein and oil-rich biomass using black soldier fly larva assisted by microbes. *J. Entomol. Zool. Stud*, 5(4), 591–597.
- Belghit, I., Liland, N. S., Gjesdal, P., Biancarosa, I., Menchetti, E., Li, Y., Waagbø, R., Kroghdahl, Å., & Lock, E.-J. (2019). Black soldier fly larvae meal can replace fish meal in diets of sea-water phase Atlantic salmon (*Salmo salar*). *Aquaculture*, 503, 609–619.
- Dayrit, F. M. (2015). The properties of lauric acid and their significance in coconut oil. *Journal of the American Oil Chemists' Society*, 92, 1–15.
- Devi, W. D., Bonysana, R., Kapesa, K., Rai, A. K., Mukherjee, P. K., & Rajashekar, Y. (2022). Potential of edible insects as source of functional foods: biotechnological approaches for improving functionality. *Systems Microbiology and Biomanufacturing*, 2(3), 461–472.
- Donadelli, V., Marco, P., Mandich, A., Finoia, M. G., Cardinaletti, G., Petochi, T., Longobardi, A., Tibaldi, E., & Marino, G. (2024). Effects of Dietary Plant Protein Replacement with Insect and Poultry By-Product Meals on the Liver Health and Serum Metabolites of Sea Bream (*Sparus aurata*) and Sea Bass (*Dicentrarchus labrax*). *Animals (Basel)*, 12;14(2):2. <https://doi.org/10.3390/ani14020241>.
- Dumas, A., Raggi, T., Barkhouse, J., Lewis, E., & Weltzien, E. (2018). The oil fraction and partially defatted meal of black soldier fly larvae (*Hermetia illucens*) affect differently growth performance, feed efficiency, nutrient deposition, blood glucose and lipid digestibility of rainbow trout (*Oncorhynchus mykiss*). *Aquaculture*, 492, 24–34.
- Gligorescu, A., Fischer, C. H., Larsen, P. F., Nørgaard, J. V., & Heckman, L.-H. L. (2020). Production and optimization of *Hermetia illucens* (L.) larvae reared on food waste and utilized as feed ingredient. *Sustainability*, 12(23), 9864.

- Heuel, M., Sandrock, C., Leiber, F., Mathys, A., Gold, M., Zurbrügg, C., Gangnat, I. D. M., Kreuzer, M., & Terranova, M. (2021). Black soldier fly larvae meal and fat can completely replace soybean cake and oil in diets for laying hens. *Poultry Science*, 100(4), 101034.
- Hossain, M. S., Small, B. C., & Hardy, R. (2023). Insect lipid in fish nutrition: Recent knowledge and future application in aquaculture. *Reviews in Aquaculture*.
- Józefiak, D., Świątkiewicz, S., Kierończyk, B., Rawski, M., Długosz, J., Engberg, R. M., & Højberg, O. (2018). Clostridium perfringens challenge and dietary fat type modifies performance, microbiota composition and histomorphology of the broiler chicken gastrointestinal tract. *Eur. Poult. Sci*, 80, 130.
- Khatun, J., Loh, T., Akit, H., Foo, H., & Mohamad, R. (2018). Influence of different sources of oil on performance, meat quality, gut morphology, ileal digestibility and serum lipid profile in broilers. *Journal of Applied Animal Research*, 46(1), 479–485.
- Kierończyk, B., Rawski, M., Józefiak, A., Mazurkiewicz, J., Świątkiewicz, S., Siwek, M., Bednarczyk, M., Szumacher-Strabel, M., Cieślak, A., & Benzertiha, A. (2018). Effects of replacing soybean oil with selected insect fats on broilers. *Animal Feed Science and Technology*, 240, 170–183.
- Kierończyk, B., Sypniewski, J., Mikołajczak, Z., Rawski, M., Pruszyńska-Oszmałek, E., Sassek, M., Kołodziejcki, P., & Józefiak, D. (2022). Replacement of soybean oil with cold-extracted fat from *Hermetia illucens* in young turkey diets: Effects on performance, nutrient digestibility, selected organ measurements, meat and liver tissue traits, intestinal microbiota modulation, and physiological. *Animal Feed Science and Technology*, 286, 115210.
- Kierończyk, B., Sypniewski, J., Rawski, M., Czeaka, W., Świątkiewicz, S., & Józefiak, D. (2020). From waste to sustainable feed material: the effect of *Hermetia illucens* oil on the growth performance, nutrient digestibility, and gastrointestinal tract morphometry of broiler chickens. *Annals of Animal Science*, 20(1), 157–177.
- Kim, B., Bang, H. T., Kim, K. H., Kim, M. J., Jeong, J. Y., Chun, J. L., & Ji, S. Y. (2020). Evaluation of black soldier fly larvae oil as a dietary fat source in broiler chicken diets. *Journal of Animal Science and Technology*, 62(2), 187.
- Kim, Y. B., Kim, D.-H., Jeong, S.-B., Lee, J.-W., Kim, T.-H., Lee, H.-G., & Lee, K.-W. (2020). Black soldier fly larvae oil as an alternative fat source in broiler nutrition. *Poultry Science*, 99(6), 3133–3143.
- Kipkoech, C., Jaster-Keller, J., Gottschalk, C., Wesonga, J., & Maul, R. (2023). African traditional use of edible insects and challenges towards the future trends of food and feed. *Journal of Insects as Food and Feed*, 1–24.
- Knarreborg, A., Simon, M. A., Engberg, R. M., Jensen, B. B., & Tannock, G. W. (2002). Effects of dietary fat source and subtherapeutic levels of antibiotic on the bacterial community in the ileum of broiler chickens at various ages. *Applied and Environmental Microbiology*, 68(12), 5918–5924.
- Kumar, V., Fawole, F. J., Romano, N., Hossain, M. S., Labh, S. N., Overturf, K., & Small, B. C. (2021). Insect (black soldier fly, *Hermetia illucens*) meal supplementation prevents the soybean meal-induced intestinal enteritis in rainbow trout and health benefits of using insect oil. *Fish & Shellfish Immunology*, 109, 116–124.
- Li, S., Ji, H., Zhang, B., Zhou, J., & Yu, H. (2017). Defatted black soldier fly (*Hermetia illucens*) larvae meal in diets for juvenile Jian carp (*Cyprinus carpio* var. Jian): Growth performance,

- antioxidant enzyme activities, digestive enzyme activities, intestine and hepatopancreas histological structure. *Aquaculture*, 477, 62–70.
- Lim, J.-W., Mohd-Noor, S.-N., Wong, C.-Y., Lam, M.-K., Goh, P.-S., Beniers, J., Oh, W.-D., Jumbri, K., & Ghani, N. A. (2019). Palatability of black soldier fly larvae in valorizing mixed waste coconut endosperm and soybean curd residue into larval lipid and protein sources. *Journal of Environmental Management*, 231, 129–136.
- Liu, Z., Minor, M., Morel, P. C., & Najar-Rodriguez, A. J. (2018). Bioconversion of three organic wastes by black soldier fly (Diptera: Stratiomyidae) larvae. *Environmental Entomology*, 47(6), 1609–1617.
- Lozica, L., Starcevic, K., Gavrilovic, A., & Masek, T. (2022). The influence of feeding media enriched with different oils on the fatty acid composition of the Black soldier fly (*Hermetia illucens*). *Vet. Arh*, 92, 291–300.
- Lu, S., Taethaisong, N., Meethip, W., Surakhunthod, J., Sinpru, B., Sroichak, T., Archa, P., Thongpea, S., Paengkoum, S., & Purba, R. A. P. (2022). Nutritional composition of black soldier fly larvae (*Hermetia illucens* L.) and its potential uses as alternative protein sources in animal diets: A review. *Insects*, 13(9), 831.
- Lumanlan, J. C., Williams, M., & Jayasena, V. (2022). Edible insects: environmentally friendly sustainable future food source. *International Journal of Food Science & Technology*, 57(10), 6317–6325.
- Nasir, N. A. M. M., Abllah, Z., Jalaludin, A. A., Shahdan, I. A., & WNH.W., A. M. (2018). Virgin coconut oil and its antimicrobial properties against pathogenic microorganisms: a review. *International Dental Conference of Sumatera Utara 2017 (IDCSU 2017)*.
- Nugroho, R., & Nur, F. (2018). Insect-based protein: future promising protein source for fish cultured. *IOP Conference Series: Earth and Environmental Science*.
- Oonincx, D. G., Broekhoven, S., Huis, A., & Loon, J. J. (2015). Feed conversion, survival and development, and composition of four insect species on diets composed of food by-products. *Plos One*, 10(12), 144601.
- Patterson, P., Acar, N., Ferguson, A., Trimble, L., Sciubba, H., & Koutsos, E. (2021). The impact of dietary Black Soldier Fly larvae oil and meal on laying hen performance and egg quality. *Poultry Science*, 100(8), 101272.
- Rawski, M., Mazurkiewicz, J., Kierończyk, B., & Józefiak, D. (2020). Black soldier fly full-fat larvae meal as an alternative to fish meal and fish oil in Siberian sturgeon nutrition: The effects on physical properties of the feed, animal growth performance, and feed acceptance and utilization. *Animals*, 10(11), 2119.
- Schiavone, A., Cullere, M., Marco, M., Meneguz, M., Biasato, I., Bergagna, S., Dezzutto, D., Gai, F., Dabbou, S., & Gasco, L. (2017). Partial or total replacement of soybean oil by black soldier fly larvae (*Hermetia illucens* L.) fat in broiler diets: Effect on growth performances, feed-choice, blood traits, carcass characteristics and meat quality. *Italian Journal of Animal Science*, 16(1), 93–100.
- Schiavone, A., Dabbou, S., Marco, M., Cullere, M., Biasato, I., Biasibetti, E., Capucchio, M. T., Bergagna, S., Dezzutto, D., & Meneguz, M. (2018). Black soldier fly larva fat inclusion in finisher broiler chicken diet as an alternative fat source. *Animal*, 12(10), 2032–2039.
- St-Hilaire, S., Cranfill, K., McGuire, M. A., Mosley, E. E., Tomberlin, J. K., Newton, L., Sealey, W., Sheppard, C., & Irving, S. (2007). Fish offal recycling by the black soldier fly produces

- a foodstuff high in omega-3 fatty acids. *Journal of the World Aquaculture Society*, 38(2), 309–313.
- Sypniewski, J., Kierończyk, B., Benzertiha, A., Mikołajczak, Z., Pruszyńska-Oszmałek, E., Kołodziejcki, P., Sassek, M., Rawski, M., Czekala, W., & Józefiak, D. (2020). Replacement of soybean oil by *Hermetia illucens* fat in turkey nutrition: Effect on performance, digestibility, microbial community, immune and physiological status and final product quality. *British Poultry Science*, 61(3), 294–302.
- Taulescu, C., Mihaiu, M., Bele, C., Matea, C., Dan, S. D., Mihaiu, R., Lapusan, A., & Ciupa, A. (2010). Manipulating the fatty acid composition of poultry meat for improving consumer's health. *Bull. UASVM*, 67(2).
- Ushakova, N., Brodskii, E., Kovalenko, A., Bastrakov, A., Kozlova, A., & D.), P. (2016). Characteristics of lipid fractions of larvae of the black soldier fly *Hermetia illucens*. *Doklady Biochemistry and Biophysics*.
- Wang, J., Wang, X., Li, J., Chen, Y., Yang, W., & Zhang, L. (2015). Effects of dietary coconut oil as a medium-chain fatty acid source on performance, carcass composition and serum lipids in male broilers. *Asian-Australasian Journal of Animal Sciences*, 28(2), 223.
- Wang, Y.-S., & Shelomi, M. (2017). Review of black soldier fly (*Hermetia illucens*) as animal feed and human food. *Foods*, 6(10), 91.
- Yoon, B. K., Jackman, J. A., Valle-González, E. R., & Cho, N.-J. (2018). Antibacterial free fatty acids and monoglycerides: biological activities, experimental testing, and therapeutic applications. *International Journal of Molecular Sciences*, 19(4).
- Zeitz, J., Fennhoff, J., Kluge, H., Stangl, G., & Eder, K. (2015). Effects of dietary fats rich in lauric and myristic acid on performance, intestinal morphology, gut microbes, and meat quality in broilers. *Poultry Science*, 94(10), 2404–2413.